

ON THE DOMINATION OF TRIANGULATED DISCS

NOOR A'LAWIAH ABD AZIZ, Penang, NADER JAFARI RAD, Tehran,
HAILIZA KAMARULHAILI, Penang

Received August 10, 2021. Published online December 5, 2022.
Communicated by Jean-Sébastien Sereni

Abstract. Let G be a 3-connected triangulated disc of order n with the boundary cycle C of the outer face of G . Tokunaga (2013) conjectured that G has a dominating set of cardinality at most $\frac{1}{4}(n+2)$. This conjecture is proved in Tokunaga (2020) for $G - C$ being a tree. In this paper we prove the above conjecture for $G - C$ being a unicyclic graph. We also deduce some bounds for the double domination number, total domination number and double total domination number in triangulated discs.

Keywords: domination; double domination; total domination; double total domination; planar graph; triangulated disc

MSC 2020: 05C69

1. INTRODUCTION

For a simple graph G with vertex set $V = V(G)$, the number of vertices of G is called the *order* of G and is denoted by $n = n(G)$. The *open neighborhood* of a vertex $v \in V$ is $N(v) = N_G(v) = \{u \in V : uv \in E\}$ and the *closed neighborhood* of v is $N[v] = N_G[v] = N(v) \cup \{v\}$. The *degree* of a vertex v , denoted by $\deg(v)$ (or $\deg_G(v)$ to refer to G), is the cardinality of its open neighborhood. For an integer $k \geq 2$, a graph G is called k -connected if it has more than k vertices and remains connected whenever fewer than k vertices are removed. The *contraction* of an edge is an operation that removes the edge while simultaneously merging the two vertices that it previously joined. A plane graph G is said to be a *triangulated disc* if it is 2-connected and all its faces are triangles except for the outer (infinite) face. The boundary cycle of the outer face of G is called the *outer cycle* of G and is denoted $C(G)$. The graph $G - C(G)$ is denoted by $\text{In}(G)$.

A subset $S \subseteq V$ is a *dominating set* of G if every vertex in $V - S$ has a neighbor in S . Note that by this definition any vertex of G dominates itself and its neighbors. The *domination number* $\gamma(G)$ is the minimum cardinality of a dominating set of G . A subset $S \subseteq V$ in a graph with no isolated vertex is a *total dominating set* of G if every vertex in V has a neighbor in S . The *total domination number* $\gamma_t(G)$ is the minimum cardinality of a total dominating set of G . For a comprehensive survey on the subject of domination parameters in graphs the reader can refer to [9].

Harary and Haynes in [8] defined a generalization of domination, namely k -tuple domination, which is called double domination if $k = 2$. A subset S of vertices of a graph G is a *double dominating set* of G if for every vertex $v \in V(G)$, $|N[v] \cap S| \geq 2$. The *double domination number* $\gamma_{\times 2}(G)$ is the minimum cardinality of a double dominating set of G . The concept of double domination in graph was studied in, for example, [2], [3], [4], [7]. A subset S of V in a graph with no isolated vertex is a *double total dominating set* of G if for every vertex $v \in V$, $|N(v) \cap S| \geq 2$. The *double total domination number* $\gamma_{\times 2,t}(G)$ is the minimum cardinality of a double total dominating set of G . The concept of double total domination was introduced in [11] and further studied in, for example, [10], [12], [16]. This concept was also studied under the name *total 2-domination*, see for example, [1], [6].

Matheson and Tarjan in [15] proved that any triangulated disc G of order n has a dominating set of cardinality at most $\frac{1}{3}n$, and conjectured that $\gamma(G) \leq \frac{1}{4}n$ for every n -vertex triangulation G with sufficiently large n . King and Pelsmajer (see [13]) proved this conjecture for graphs of maximum degree 6. Campos and Wakabayashi in [5] and Tokunaga in [17] independently proved that $\gamma(G) \leq \frac{1}{4}(n + t)$ for each n -vertex outerplanar graph G of order n having t vertices of degree two. An improvement of the $\frac{1}{4}(n + t)$ -bound is given by Li et al. in [14]. Tokunaga (see [17]) also posed the following conjecture.

Conjecture 1 (Tokunaga, [17]). If G is a 3-connected n -vertex triangulated disc, then $\gamma(G) \leq \frac{1}{4}(n + 2)$.

Recently, Tokunaga in [18] proved Conjecture 1 for triangulated discs G such that $\text{In}(G)$ is a tree.

Theorem 2 (Tokunaga, [18]). If G is an n -vertex triangulated disc such that $\text{In}(G)$ is a tree and $C(G)$ is an induced cycle of G , then $\gamma(G) \leq \frac{1}{4}(n + 2)$.

In this paper we proved a stronger version of Conjecture 1 for triangulated discs G such that $\text{In}(G)$ is a unicyclic graph. We show that if G is an n -vertex triangulated disc such that $\text{In}(G)$ is a unicyclic graph and $C(G)$ is an induced cycle of G , then $\gamma(G) \leq \frac{1}{4}(n + 1)$. We also apply the methods for double domination, total domination and double total domination numbers in triangulated discs.

We use the same method given in [18] using proper colorings. For a given integer $k \geq 1$, a function $f: V(G) \rightarrow \{1, \dots, k\}$ is called a *proper k -coloring* if $f(u) \neq f(v)$ for every edge uv of G . If f is proper k -coloring and a vertex v is dominated by a vertex of color i for some $i \in \{1, \dots, k\}$, then we say that v is *dominated by color i* . We use the following key lemma of [18].

Lemma 3 (Tokunaga, [18]). *Let G be an n -vertex triangulated disc such that $\text{In}(G)$ is a tree and $C(G)$ is an induced cycle of G , and let v be a vertex of $C(G)$ with $\deg_G(v) = 3$. Then $G - v$ has a proper 4-coloring such that each vertex of $G - v$ is dominated by all the four colors except the vertices of $N_G(v)$.*

2. BOUNDS

We first present a bound for the domination number.

Theorem 4. *If G is an n -vertex triangulated disc such that $\text{In}(G)$ is a unicyclic graph and $C(G)$ is an induced cycle of G , then $\gamma(G) \leq \frac{1}{4}(n+1)$. This bound is sharp.*

Proof. Let G be an n -vertex triangulated disc such that $\text{In}(G)$ is a unicyclic graph and $C(G)$ is an induced cycle of G . Clearly, $\text{In}(G)$ contains a triangle, since all faces are triangles except for the outer (infinite) face. Let abc be a triangle in $\text{In}(G)$, and G^* be the graph obtained from G by contraction of the edge ab . Then G^* is an $(n - 1)$ -vertex triangulated disc such that $\text{In}(G)$ is a tree, namely T . Clearly, $C(G^*) = C(G)$ and $\{a, b, c\} \cap V(C) = \emptyset$. Let v be a vertex of $C(G^*)$ with $\deg_{G^*}(v) = 3$. Clearly, $\deg_{G^*}(v) = \deg_G(v) = 3$. Let $N_{G^*}(v) = \{u, w, x\}$, where $N_C(v) = \{u, w\}$ and x is the unique vertex of T , which is adjacent to v in G^* . Now we follow the proof of Theorem 2 given in [18]. By Lemma 3, $G^* - v$ has a proper 4-coloring f such that each vertex of $G^* - v$ is dominated by all the four colors except the vertices of $N_{G^*}(v)$. Let G' be the $(n + 1)$ -vertex graph such that $V(G') = V(G^*) \cup \{p, q\}$ and $E(G') = E(G^*) \cup \{pu, pv, pw, qu, qv, qw\}$.

We define a 4-coloring f' on G^* as follows. Define $f'(y) = f(y)$ if $y \in V(G^*) - \{v\}$, and let $f'(v)$ be a color different from $f'(x)$, $f'(u)$ and $f'(w)$. Now assign $f'(p)$ and $f'(q)$ such that $\{f'(x), f'(p), f'(q), f'(v)\} = \{1, 2, 3, 4\}$. Then each vertex of G^* is dominated by all the four colors. By renaming the colors, if necessary, we may assume that $|\{y \in V(G^*): f'(y) = f'(c) = f(c)\}| \leq \frac{1}{4}(n + 1)$. Let $S = \{y \in V(G^*): f'(y) = f'(c) = f(c)\}$. Now, we form a set S' defined by $S' = S$ if $S \cap \{p, q\} = \emptyset$, $S' = (S - \{p\}) \cup \{v\}$ if $p \in S$, and $S' = (S - \{q\}) \cup \{v\}$ if $q \in S$. Then S' is a dominating set for G^* of cardinality at most $\frac{1}{4}(n + 1)$. Since $c \in S'$, we deduce that S' is a dominating set for G , and the proof is completed.




Figure 1. The graph H .

For the sharpness, see the graph H depicted in Figure 1. It is easy to see that the set $\{u_5, u_8, u_{10}, u_{15}\}$ is a minimum dominating set. \square

Next apply the above arguments for double domination, total domination, and double total domination. Following the proofs of Theorem 2, we can assume that $|\{y: f'(y) \in \{1, 2\}\}| \leq \frac{1}{2}(n + 2)$. Then we form a double dominating set for G of cardinality at most $\frac{1}{2}(n + 2)$. Thus, we have the following.

Corollary 5. *If G is an n -vertex triangulated disc such that $\text{In}(G)$ is a tree and $C(G)$ is an induced cycle of G , then $\gamma_{\times 2}(G) \leq \frac{1}{2}(n + 2)$.*

We next prove an upper bound for the double domination number for G when $\text{In}(G)$ is a unicyclic graph.

Theorem 6. *If G is an n -vertex triangulated disc such that $\text{In}(G)$ is a unicyclic graph and $C(G)$ is an induced cycle of G , then $\gamma_{\times 2}(G) \leq \frac{1}{2}(n + 1)$.*

PROOF. We follow the proof of Theorem 4. Let G be an n -vertex triangulated disc such that $\text{In}(G)$ is a unicyclic graph and $C(G)$ is an induced cycle of G . Let abc be a triangle in $\text{In}(G)$, and G^* be the graph obtained from G by contraction of the edge ab . Then G^* is an $(n - 1)$ -vertex triangulated disc such that $\text{In}(G)$ is a tree. Clearly, a and b have a common neighbor $d \neq c$ in G , since G is a triangulated disc. Now, following the proof, we may assume that $|\{y: f'(y) \in \{f'(c), f'(d)\}\}| \leq \frac{1}{2}(n + 1)$. Let $S = \{y: f'(y) \in \{f'(c), f'(d)\}\}$, and form S' as described in the proof of Theorem 4. Since $c, d \in S'$, we obtain that S' is a dominating set for G , and the proof is completed. \square

It is evident that $\gamma_t(G) \leq \gamma_{\times 2}(G)$ for any graph G with no isolated vertex. Thus, the bounds given in Corollary 5 and Theorem 6 are also valid for total domination. We next prove upper bounds for the double total domination number.

Theorem 7. *Let G be an n -vertex triangulated disc such that $C(G)$ is an induced cycle of G . If $\text{In}(G)$ is a tree, then $\gamma_{\times 2, t}(G) \leq \frac{3}{4}(n + 2)$, and if $\text{In}(G)$ is a unicyclic graph, then $\gamma_{\times 2, t}(G) \leq \frac{3}{4}(n + 1)$.*

Proof. We follow the proof of Theorems 2 and 4. First assume that $\text{In}(G)$ is a tree. Let f' be the given 4-coloring in the proof of Theorem 2, and assume that $|\{y: f'(y) \neq 1\}| \leq \frac{3}{4}(n+2)$. Let $S = \{y: f'(y) \neq 1\}$ and S' be formed from S as described in the proof of Theorem 2. Then S' is a double total dominating set for G of cardinality at most $\frac{3}{4}(n+2)$. Next assume that $\text{In}(G)$ is a unicyclic graph. We follow the proof of Theorem 4. Let y^* be the vertex formed by contraction of the edge ab . Let f' be the given 4-coloring in the proof of Theorem 4, and assume that $|\{y: f'(y) \neq f'(y^*)\}| \leq \frac{3}{4}(n+1)$. Let $S = \{y: f'(y) \neq f'(y^*)\}$ and S' be formed from S as described in the proof of Theorem 4. Then S' is a double total dominating set for G of cardinality at most $\frac{3}{4}(n+1)$. \square

We close with the following conjecture.

Conjecture 8. If G is an n -vertex triangulated disc, then $\gamma(G) \leq \frac{1}{4}(n+2-t)$, $\gamma_{\times 2}(G) \leq \frac{1}{2}(n+2-t)$ and $\gamma_{\times 2,t}(G) \leq \frac{3}{4}(n+2-t)$, where t is the number of vertex-disjoint triangles in $\text{In}(G)$.

References

- [1] *S. Bermudo, J. L. Sánchez, J. M. Sigarreta*: Total k -domination in Cartesian product graphs. *Period. Math. Hung.* *75* (2017), 255–267. [zbl](#) [MR](#) [doi](#)
- [2] *M. Blidia, M. Chellali, T. W. Haynes*: Characterizations of trees with equal paired and double domination numbers. *Discrete Math.* *306* (2006), 1840–1845. [zbl](#) [MR](#) [doi](#)
- [3] *A. Cabrera-Martínez*: New bounds on the double domination number of trees. *Discrete Appl. Math.* *315* (2022), 97–103. [zbl](#) [MR](#) [doi](#)
- [4] *A. Cabrera-Martínez, J. A. Rodríguez-Velázquez*: A note on double domination in graphs. *Discrete Appl. Math.* *300* (2021), 107–111. [zbl](#) [MR](#) [doi](#)
- [5] *C. N. Campos, Y. Wakabayashi*: On dominating sets of maximal outerplanar graphs. *Discrete Appl. Math.* *161* (2013), 330–335. [zbl](#) [MR](#) [doi](#)
- [6] *H. Fernau, J. A. Rodríguez-Velázquez, J. M. Sigarreta*: Global powerful r -alliances and total k -domination in graphs. *Util. Math.* *98* (2015), 127–147. [zbl](#) [MR](#)
- [7] *J. Harant, M. A. Henning*: On double domination in graphs. *Discuss. Math., Graph Theory* *25* (2005), 29–34. [zbl](#) [MR](#) [doi](#)
- [8] *F. Harary, T. W. Haynes*: Double domination in graphs. *Ars Comb.* *55* (2000), 201–213. [zbl](#) [MR](#)
- [9] *T. W. Haynes, S. T. Hedetniemi, P. J. Slater*: *Fundamentals of Domination in Graphs*. Pure and Applied Mathematics, Marcel Dekker 208. Marcel Dekker, New York, 1998. [zbl](#) [MR](#) [doi](#)
- [10] *M. A. Henning, N. Jafari Rad*: Upper bounds on the k -tuple (Roman) domination number of a graph. *Graphs Comb.* *37* (2021), 325–336. [zbl](#) [MR](#) [doi](#)
- [11] *M. A. Henning, A. P. Kazemi*: k -tuple total domination in graphs. *Discrete Appl. Math.* *158* (2010), 1006–1011. [zbl](#) [MR](#) [doi](#)
- [12] *M. A. Henning, A. Yeo*: Strong transversals in hypergraphs and double total domination in graphs. *SIAM J. Discrete Math.* *24* (2010), 1336–1355. [zbl](#) [MR](#) [doi](#)
- [13] *E. L. C. King, M. J. Pelsmayer*: Dominating sets in plane triangulations. *Discrete Math.* *310* (2010), 2221–2230. [zbl](#) [MR](#) [doi](#)
- [14] *Z. Li, E. Zhu, Z. Shao, J. Xu*: On dominating sets of maximal outerplanar and planar graphs. *Discrete Appl. Math.* *198* (2016), 164–169. [zbl](#) [MR](#) [doi](#)

- [15] *L. R. Matheson, R. E. Tarjan*: Dominating sets in planar graphs. *Eur. J. Comb.* *17* (1996), 565–568. [zbl](#) [MR](#) [doi](#)
- [16] *D. Pradhan*: Algorithmic aspects of k -tuple total domination in graphs. *Inf. Process. Lett.* *112* (2012), 816–822. [zbl](#) [MR](#) [doi](#)
- [17] *S. Tokunaga*: Dominating sets of maximal outerplanar graphs. *Discrete Appl. Math.* *161* (2013), 3097–3099. [zbl](#) [MR](#) [doi](#)
- [18] *S. Tokunaga*: On domination number of triangulated discs. *J. Inf. Process.* *28* (2020), 846–848. [doi](#)

Authors' addresses: *Noor A'lawiah Abd Aziz*, School of Mathematical Sciences, Universiti Sains Malaysia, 11800 USM Penang, Malaysia, e-mail: nooralawiah@gmail.com; *Nader Jafari Rad* (corresponding author), Department of Mathematics, Shahed University, Tehran, Iran, e-mail: n.jafarirad@gmail.com; *Hailiza Kamarulhaili*, School of Mathematical Sciences, Universiti Sains Malaysia, 11800 USM Penang, Malaysia, e-mail: hailiza@usm.my.