

EXISTENCE, UNIQUENESS AND CONTINUITY RESULTS  
OF WEAK SOLUTIONS FOR NONLOCAL NONLINEAR  
PARABOLIC PROBLEMS

TAYEB BENHAMOUD, ELMEHDI ZAOUCHÉ, El Oued,  
MAHMOUD BOUSSELSAL, Algiers

Received April 26, 2023. Published online February 29, 2024.  
Communicated by Ondřej Kreml

*Abstract.* This paper is concerned with the study of a nonlocal nonlinear parabolic problem associated with the equation  $u_t - M(\int_{\Omega} \phi u \, dx) \operatorname{div} (A(x, t, u) \nabla u) = g(x, t, u)$  in  $\Omega \times (0, T)$ , where  $\Omega$  is a bounded domain of  $\mathbb{R}^n$  ( $n \geq 1$ ),  $T > 0$  is a positive number,  $A(x, t, u)$  is an  $n \times n$  matrix of variable coefficients depending on  $u$  and  $M: \mathbb{R} \rightarrow \mathbb{R}$ ,  $\phi: \Omega \rightarrow \mathbb{R}$ ,  $g: \Omega \times (0, T) \times \mathbb{R} \rightarrow \mathbb{R}$  are given functions. We consider two different assumptions on  $g$ . The existence of a weak solution for this problem is proved using the Schauder fixed point theorem for each of these assumptions. Moreover, if  $A(x, t, u) = a(x, t)$  depends only on the variable  $(x, t)$ , we investigate two uniqueness theorems and give a continuity result depending on the initial data.

*Keywords:* nonlocal nonlinear parabolic problem; Schauder fixed point theorem; weak solution; existence; uniqueness

*MSC 2020:* 35D30, 35K55, 35Q92

## 1. INTRODUCTION

Let  $\Omega$  be a bounded domain in  $\mathbb{R}^n$  ( $n \geq 1$ ),  $T$  be a positive number,  $Q = \Omega \times (0, T)$  and  $\phi, u_0 \in L^2(\Omega)$ . Let  $A: Q \times \mathbb{R} \rightarrow \mathbb{R}^{n \times n}$  be a vector function such that  $(x, t) \mapsto A(x, t, s)$  is measurable for all  $s \in \mathbb{R}$ ,  $s \mapsto A(x, t, s)$  is continuous for a.e.  $(x, t) \in Q$  and

$$(1.1) \quad \forall \xi \in \mathbb{R}^n: \lambda |\xi|^2 \leq A(x, t, s) \xi \cdot \xi,$$

$$(1.2) \quad \forall \xi \in \mathbb{R}^n: |A(x, t, s) \xi| \leq \Lambda |\xi|,$$

for all  $s \in \mathbb{R}$  and a.e.  $(x, t) \in Q$ , where  $\lambda$  and  $\Lambda$  are positive constants, and let  $M: \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function satisfying for some constants  $m_1 \geq m_0 > 0$ ,

$$(1.3) \quad m_0 \leq M(s) \leq m_1 \quad \forall s \in \mathbb{R}.$$

On the other hand, let  $g: Q \times \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $(x, t) \mapsto g(x, t, s)$  is measurable for all  $s \in \mathbb{R}$ ,  $g(\cdot, \cdot, 0) \in L^2(Q)$  and satisfies one of the two following assumptions:

$$(1.4) \quad \forall s_1, s_2 \in \mathbb{R}, \text{ a.e. } (x, t) \in Q:$$

$$|g(x, t, s_1) - g(x, t, s_2)| \leq \begin{cases} b_p(x, t)|s_1 - s_2|^p, & p \in (0, 1); \\ b_1(x, t)|s_1 - s_2|, & p = 1, \end{cases}$$

where  $b_p \in L^{2/(1-p)}(Q)$  for  $p \in (0, 1)$ ,  $b_1 \in L^\infty(Q)$ ,  $C_\Omega(2T/(\lambda m_0))^{1/2} \|b_1\|_{L^\infty(Q)} < 1$  with  $C_\Omega$  denoting a Poincaré constant for  $H_0^1(\Omega)$ , and for a.e.  $(x, t) \in Q$ ,  $s \mapsto g(x, t, s)$  is continuous,

$$(1.5) \quad \exists h \in L^2(Q) \quad \forall s \in \mathbb{R}, \text{ a.e. } (x, t) \in Q: |g(x, t, s)| \leq h(x, t).$$

We now consider the following weak formulation of nonlocal nonlinear parabolic problems:

$$(1.6) \quad \begin{cases} u \in L^2(0, T; H_0^1(\Omega)) \cap C([0, T]; L^2(\Omega)), & u_t \in L^2(0, T; H^{-1}(\Omega)), \\ u(\cdot, 0) = u_0 \quad \text{a.e. in } \Omega, \\ \frac{d}{dt}(u, \xi) + M(l(u)) \int_\Omega A(x, t, u) \nabla u \cdot \nabla \xi \, dx = \int_\Omega g(x, t, u) \xi \, dx \\ \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega), \end{cases}$$

where

$$(u, \xi) = \int_\Omega u \xi \, dx \quad \text{and} \quad l(u) = l(u)(t) = \int_\Omega \phi(x) u(x, t) \, dx.$$

The study of nonlocal problems has attracted the attention of many authors and several results have been established. These types of problems (1.6) arise in a wide variety of applications in physics and population dynamics. For instance, the solution  $u$  can be used to describe the population density of bacteria in space and time. Also, this model can address some questions concerning the heat conduction. See [7], [8], [9], [10], [17] for more details.

In the homogenous case when  $A(x, t, u)$  is the identity matrix, the authors in [8] and [3] established existence theorems for (1.6) by using the Galerkin Method, respectively, for the second term  $g$  depending only on  $(x, t)$ ,  $g(x, t, u) = g(x, t) \in$

$L^2(0, T; H^{-1}(\Omega))$  and for  $g(x, t, u) = k(x) - f(u)$  such that  $k \in L^2(\Omega)$  and  $f$  is a continuously differentiable function satisfying  $f(u)u \geq -\mu u^2 - c_1$ ,  $f'(u) \geq -\alpha$ , where  $c_1, \alpha$  are two positive constants,  $0 < \mu < m_0 \lambda_1$  with  $\lambda_1 > 0$  is the first eigenvalue of the operator  $(-\Delta, H_0^1(\Omega))$ . Moreover, the uniqueness and the asymptotic behavior of solutions and other results have been obtained under some additional conditions. By means of the Schauder fixed point theorem, the existence of weak solutions for (1.6) has been proved in [13] for a Lipschitz continuous function  $g(x, t, u) = f(u)$  such that  $f(0) = 0$  and  $f'(0)$  exists. Also, the uniqueness and the result on existence of periodic solution have been established. If  $g$  depends only on the variable  $(x, t)$ ,  $g(x, t, u) = g(x, t) \in L^2(0, T; H^{-1}(\Omega))$  and  $l: L^2(\Omega) \rightarrow \mathbb{R}$  is a continuous function, the authors in [10] gave an existence theorem for (1.6) and by imposing additional conditions, they investigated the uniqueness and asymptotic behavior of solutions. In [16], the authors considered the following nonlocal parabolic problem:

$$\begin{cases} u_t - a\left(\int_{\Omega} |u|^{\gamma} dx\right) \Delta u = f(u) & \text{in } \Omega \times (0, T), \\ u = 0 & \text{on } \partial\Omega \times (0, T), \\ u(\cdot, 0) = u_0 & \text{in } \Omega, \end{cases}$$

where  $\Omega$  is a sufficiently regular domain,  $\gamma \in [1, \infty)$ ,  $T \in (0, \infty]$ ,  $u_0 \in C^{2+\alpha}(\bar{\Omega})$ ,  $\alpha \in (0, 1)$ ,  $f \in C^1(\mathbb{R})$  and  $a \in C^1([0, \infty))$  with  $\inf_{t \in [0, \infty)} a(t) \geq a(0) := a_0 > 0$ .

By using the sub-supersolution method, they proved the existence, uniqueness and long-time behavior of positive solutions.

For the nonlocal problems in the stationary case, we refer to [1], [2], [4], [5], [9], [11], [15], [18], [19], [20] and the references therein in which various methods have been used for studying the existence and uniqueness of solutions and other questions.

In this paper, we suppose that (1.1)–(1.3) and one of (1.4) and (1.5) hold. We prove the existence of a solution for each problem (1.6) by applying the Schauder fixed point theorem. Moreover, if  $A(x, t, u) = a(x, t)$  depends only on the variable  $(x, t)$ , we investigate two uniqueness theorems and give a continuity result depending on the initial data as in [10], Lemma 5.1, assuming that  $M$  is a Lipschitz continuous function. Our first uniqueness theorem is concerned with the case  $p = 1$  in which the function  $g$  satisfies a generalized Lipschitz condition and under the assumption that  $g$  is decreasing with respect to  $u$ , we state and prove our second uniqueness theorem.

## 2. EXISTENCE OF A SOLUTION

In this section, we prove the existence of a solution of (1.6) assuming that the function  $g$  satisfies one of assumptions (1.4) and (1.5).

**Theorem 2.1.** *If (1.1)–(1.3) and either (1.4) or (1.5) holds, problem (1.6) has a solution.*

*Proof.* We apply the Schauder fixed point theorem to show the existence of a solution of (1.6). For a fixed element  $v$  of  $L^2(Q)$  and for a.e.  $t \in (0, T)$ , we define

$$\begin{aligned} A_v(u, \xi, t) &= M(l(v)) \int_{\Omega} A(x, t, v) \nabla u \cdot \nabla \xi \, dx \quad \forall u, \xi \in H_0^1(\Omega), \\ f_v(x, t) &= g(x, t, v) \quad \text{a.e. } x \in \Omega. \end{aligned}$$

Observe that  $A_v$  is a bilinear form on  $H_0^1(\Omega) \times H_0^1(\Omega)$  which satisfies

$$t \mapsto A_v(u, \xi, t) \text{ is measurable } \forall u, \xi \in H_0^1(\Omega),$$

and by (1.1)–(1.3) and the Cauchy-Schwarz inequality, we have for a.e.  $t \in (0, T)$ ,

$$\begin{aligned} |A_v(u, \xi, t)| &\leq \Lambda m_1 \|u\|_{H_0^1(\Omega)} \|\xi\|_{H_0^1(\Omega)} \quad \forall u, \xi \in H_0^1(\Omega), \\ A_v(u, u, t) + \lambda m_0 \|u\|_{L^2(\Omega)}^2 &\geq \lambda m_0 \|u\|_{H_0^1(\Omega)}^2 \quad \forall u \in H_0^1(\Omega). \end{aligned}$$

On the other hand, if one of assumptions (1.4) and (1.5) holds, the function  $f_v$  belongs to  $L^2(Q)$ . Thus, from [6], Theorem 11.7, we deduce that for each  $v \in L^2(Q)$ , there exists a unique solution of the following problem:

$$(2.1) \quad \begin{cases} u \in L^2(0, T; H_0^1(\Omega)), & u_t \in L^2(0, T; H^{-1}(\Omega)), \\ u(\cdot, 0) = u_0 & \text{a.e. in } \Omega, \\ \frac{d}{dt}(u, \xi) + M(l(v)) \int_{\Omega} A(x, t, v) \nabla u \cdot \nabla \xi \, dx = \int_{\Omega} g(x, t, v) \xi \, dx \\ & \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega). \end{cases}$$

Let us now define  $F: L^2(Q) \rightarrow L^2(0, T; H_0^1(\Omega))$  by  $F(v) = u$ . If we choose  $\xi = u$  as a test function in (2.1), we obtain

$$(2.2) \quad \frac{1}{2} \frac{d}{dt} \|u\|_{L^2(\Omega)}^2 + M(l(v)) \int_{\Omega} A(x, t, v) \nabla u \cdot \nabla u \, dx = \int_{\Omega} g(x, t, v) u \, dx.$$

Integrating (2.2) from 0 to  $t$ , we get

$$(2.3) \quad \begin{aligned} \frac{1}{2} \|u\|_{L^2(\Omega)}^2 + \int_0^t M(l(v)) \int_{\Omega} A(x, s, v) \nabla u \cdot \nabla u \, dx \, ds \\ = \int_0^t \int_{\Omega} g(x, s, v) u \, dx \, ds + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2. \end{aligned}$$

Using (1.1), (1.3), (1.4), Hölder's inequality and Poincaré's inequality, we obtain from (2.3) for all  $p \in (0, 1)$ ,

$$\begin{aligned}
& \frac{1}{2} \|u\|_{L^2(\Omega)}^2 + \lambda m_0 \int_0^t \|u\|_{H_0^1(\Omega)}^2 \, ds \\
& \leq \int_0^t \int_{\Omega} (b_p(x, s)|v|^p + |g(x, s, 0)|)|u| \, dx \, ds + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2 \\
& \leq (\|b_p\|_{L^{2/(1-p)}(Q)} \|v\|_{L^2(Q)}^p + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) \|u\|_{L^2(\Omega \times (0, t))} + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2 \\
& \leq C_{\Omega} (\|b_p\|_{L^{2/(1-p)}(Q)} \|v\|_{L^2(Q)}^p + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) \left( \int_0^t \|u\|_{H_0^1(\Omega)}^2 \, ds \right)^{1/2} + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2,
\end{aligned}$$

which, by using Young's inequality, leads to

$$\begin{aligned}
(2.4) \quad & \frac{1}{2} \|u\|_{L^2(\Omega)}^2 + \frac{\lambda m_0}{2} \int_0^t \|u\|_{H_0^1(\Omega)}^2 \, ds \\
& \leq \frac{C_{\Omega}^2}{2\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)} \|v\|_{L^2(Q)}^p + \|g(\cdot, \cdot, 0)\|_{L^2(Q)})^2 + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2 \\
& \leq \frac{C_{\Omega}^2}{\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)}^2 \|v\|_{L^2(Q)}^{2p} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2.
\end{aligned}$$

From (2.4), we derive the following estimates:

$$\begin{aligned}
(2.5) \quad & \|u\|_{L^2(0, T; H_0^1(\Omega))} \\
& \leq \frac{1}{(\lambda m_0)^{1/2}} \left\{ \frac{2C_{\Omega}^2}{\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)}^2 \|v\|_{L^2(Q)}^{2p} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \|u_0\|_{L^2(\Omega)}^2 \right\}^{1/2}
\end{aligned}$$

and

$$\begin{aligned}
(2.6) \quad & \|u\|_{L^2(Q)} \\
& \leq T^{1/2} \left\{ \frac{2C_{\Omega}^2}{\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)}^2 \|v\|_{L^2(Q)}^{2p} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \|u_0\|_{L^2(\Omega)}^2 \right\}^{1/2} \\
& \leq T^{1/2} \left\{ C_{\Omega} \left( \frac{2}{\lambda m_0} \right)^{1/2} (\|b_p\|_{L^{2/(1-p)}(Q)} \|v\|_{L^2(Q)}^p + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) + \|u_0\|_{L^2(\Omega)} \right\}.
\end{aligned}$$

Moreover, if  $p = 1$ , we arrive at

$$\begin{aligned}
(2.7) \quad & \|u\|_{L^2(0, T; H_0^1(\Omega))} \\
& \leq \frac{1}{(\lambda m_0)^{1/2}} \left\{ \frac{2C_{\Omega}^2}{\lambda m_0} (\|b_1\|_{L^{\infty}(Q)}^2 \|v\|_{L^2(Q)}^2 + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \|u_0\|_{L^2(\Omega)}^2 \right\}^{1/2}
\end{aligned}$$

and

$$(2.8) \quad \|u\|_{L^2(Q)} \leq T^{1/2} \left\{ C_{\Omega} \left( \frac{2}{\lambda m_0} \right)^{1/2} (\|b_1\|_{L^{\infty}(Q)} \|v\|_{L^2(Q)} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) + \|u_0\|_{L^2(\Omega)} \right\}.$$

By using (2.5)–(2.8) and the fact that  $C_\Omega(2T/(\lambda m_0))^{1/2}\|b_1\|_{L^\infty(Q)} < 1$ , we deduce that for all  $p \in (0, 1]$ , there exist positive constants  $R_1 = R_1(p)$ ,  $R_2 = R_2(p)$  such that

$$(2.9) \quad \|v\|_{L^2(Q)} \leq R_1 \Rightarrow (\|u\|_{L^2(Q)} \leq R_1 \text{ and } \|u\|_{L^2(0,T;H_0^1(\Omega))} \leq R_2).$$

In particular,  $F(\overline{B}(0, R_1)) \subset \overline{B}(0, R_1)$ , where  $\overline{B}(0, R_1)$  denotes the closed ball in  $L^2(Q)$  of center 0 and radius  $R_1$ . Also, if  $g$  satisfies (1.5), we can find another positive constant  $R_3$  independent of  $v$  such that  $F$  sends  $L^2(Q)$  to  $\overline{B}(0, R_3)$  and

$$(2.10) \quad \|u\|_{L^2(0,T;H_0^1(\Omega))} \leq R_3.$$

Now, we prove that  $F: \overline{B}(0, R_4) \rightarrow \overline{B}(0, R_4)$  is continuous, where  $R_4 = R_1$  or  $R_3$ . Let  $(v_j)_{j \in \mathbb{N}}$  be a sequence in  $\overline{B}(0, R_4)$  which converges to  $v_\infty \in \overline{B}(0, R_4)$  and set  $u_j = F(v_j)$ ,  $u_\infty = F(v_\infty)$ . There exists a subsequence  $j_k$  such that

$$(2.11) \quad v_{j_k} \rightarrow v_\infty \quad \text{strongly in } L^2(Q),$$

$$(2.12) \quad v_{j_k} \rightarrow v_\infty \quad \text{a.e. in } Q,$$

$$(2.13) \quad l(v_{j_k}) \rightarrow l(v_\infty) \quad \text{strongly in } L^2(0, T).$$

The last equation of (2.1) can be written in  $L^2(0, T; H^{-1}(\Omega))$  as

$$(2.14) \quad \frac{du}{dt} - M(l(v)) \operatorname{div}(A(x, t, v) \nabla u) = g(x, t, v).$$

Writing (2.14) for  $u = u_{j_k}$  and  $v = v_{j_k}$ , and multiplying the obtained equation by  $\xi \in \mathcal{D}(0, T; H_0^1(\Omega))$ , we get

$$(2.15) \quad \int_Q u_{j_k} \xi_t \, dx \, dt = \int_0^T M(l(v_{j_k})) \int_\Omega A(x, t, v_{j_k}) \nabla u_{j_k} \cdot \nabla \xi \, dx \, dt \\ - \int_Q g(x, t, v_{j_k}) \xi \, dx \, dt.$$

Using (1.2), (1.3), the Cauchy-Schwarz inequality, Poincaré's inequality and either (1.4) with (2.9) or (1.5) with (2.10), we obtain from (2.15) for a constant  $c_2$  independent of  $j_k$  and  $\xi$ ,

$$\forall k \in \mathbb{N}: \left| \int_Q u_{j_k} \xi_t \, dx \, dt \right| \leq c_2 \|\xi\|_{L^2(0,T;H_0^1(\Omega))},$$

which means that

$$(2.16) \quad \forall k \in \mathbb{N}: \|u_{j_k}\|_{L^2(0,T;H^{-1}(\Omega))} \leq c_2.$$

We notice that the balls of  $E = \{v \in L^2(0, T; H_0^1(\Omega)) / v_t \in L^2(0, T; H^{-1}(\Omega))\}$ , which is a Hilbert space when equipped with the norm

$$\{\|v\|_{L^2(0, T; H_0^1(\Omega))}^2 + \|v_t\|_{L^2(0, T; H^{-1}(\Omega))}^2\}^{1/2}$$

(see [6]), are relatively compact in  $L^2(0, T; L^2(\Omega)) = L^2(Q)$  (see [12]). Then, by (2.9), (2.10) and (2.16), the set  $F(\overline{B}(0, R_4))$  is relatively compact in  $\overline{B}(0, R_4)$ . On the other hand, there exist a subsequence, which we still denote by  $j_k$ , and  $\overline{u} \in L^2(0, T; H_0^1(\Omega))$  such that

$$(2.17) \quad u_{j_k} \rightharpoonup \overline{u} \quad \text{weakly in } L^2(0, T; H_0^1(\Omega)),$$

$$(2.18) \quad u_{j_k t} \rightharpoonup \overline{u}_t \quad \text{weakly in } L^2(0, T; H^{-1}(\Omega)),$$

$$(2.19) \quad u_{j_k} \rightarrow \overline{u} \quad \text{strongly in } L^2(Q).$$

For  $\varphi \in \mathcal{D}(0, T)$  and  $\xi \in H_0^1(\Omega)$  we have

$$(2.20) \quad \int_0^T \varphi(t) M(l(v_{j_k})) \int_{\Omega} A(x, t, v_{j_k}) \nabla u_{j_k} \cdot \nabla \xi \, dx \, dt - \int_Q u_{j_k} \xi \varphi_t \, dx \, dt \\ = \int_Q g(x, t, v_{j_k}) \xi \varphi \, dx \, dt.$$

Observe that from (1.2), (1.3), (2.12), (2.13), (2.17), the continuity of  $M$  and  $s \mapsto A(\cdot, \cdot, s)$ , and the dominated convergence theorem, we have up to a subsequence

$$(2.21) \quad M(l(v_{j_k})) \nabla \xi \rightarrow M(l(v_{\infty})) \nabla \xi \quad \text{strongly in } (L^2(Q))^n,$$

$$(2.22) \quad A(\cdot, \cdot, v_{j_k}) \nabla u_{j_k} \rightharpoonup A(\cdot, \cdot, v_{\infty}) \nabla \overline{u} \quad \text{weakly in } (L^2(Q))^n.$$

On the other hand, if (1.4) or (1.5) holds, then

$$(2.23) \quad \lim_{k \rightarrow \infty} \int_Q g(x, t, v_{j_k}) \xi \, dx \, dt = \int_Q g(x, t, v_{\infty}) \xi \, dx \, dt.$$

Indeed, if (1.4) is satisfied, Hölder's inequality gives

$$\left| \int_Q (g(x, t, v_{j_k}) - g(x, t, v_{\infty})) \xi \varphi \, dx \, dt \right| \\ \leq \begin{cases} \|b_p\|_{L^{2/(1-p)}(Q)} \|v_{j_k} - v_{\infty}\|_{L^2(Q)}^p \|\xi \varphi\|_{L^2(Q)}, & p \in (0, 1); \\ \|b_1\|_{L^{\infty}(Q)} \|v_{j_k} - v_{\infty}\|_{L^2(Q)} \|\xi \varphi\|_{L^2(Q)}, & p = 1, \end{cases}$$

which leads to (2.23) by passing to the limit as  $k \rightarrow \infty$  and using (2.11). Also, if  $g$  satisfies (1.5), it is sufficient to use (2.12), the continuity of  $s \mapsto g(\cdot, \cdot, s)$  and the

dominated convergence theorem to get (2.23). Then, passing to the limit in (2.20) as  $k \rightarrow \infty$  and using (2.19) and (2.21)–(2.23), we obtain

$$\int_0^T \varphi(t) M(l(v_\infty)) \int_\Omega A(x, t, v_\infty) \nabla \bar{u} \cdot \nabla \xi \, dx \, dt - \int_Q \bar{u} \xi \varphi_t \, dx \, dt = \int_Q g(x, t, v_\infty) \xi \varphi \, dx \, dt,$$

which can be written as

$$\begin{aligned} & \frac{d}{dt} (\bar{u}, \xi) + M(l(v_\infty)) \int_\Omega A(x, t, v_\infty) \nabla \bar{u} \cdot \nabla \xi \, dx \\ &= \int_\Omega g(x, t, v_\infty) \xi \, dx \quad \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega). \end{aligned}$$

Let us prove that  $\bar{u}(\cdot, 0) = u_0$  a.e. in  $\Omega$ . For a.e.  $t \in (0, T)$  and  $\xi \in H_0^1(\Omega)$ , we have

$$(2.24) \quad \int_0^t \langle u_{j_k t}, \xi \rangle \, ds = (u_{j_k}(\cdot, t), \xi) - (u_{j_k}(\cdot, 0), \xi) = (u_{j_k}(\cdot, t), \xi) - (u_0, \xi),$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality product between  $H^{-1}(\Omega)$  and  $H_0^1(\Omega)$ . We notice that from (2.19) we have up to a subsequence

$$(2.25) \quad u_{j_k} \rightarrow \bar{u} \quad \text{strongly in } L^2(\Omega), \text{ a.e. in } (0, T),$$

then, passing to the limit in (2.24) as  $k \rightarrow \infty$ , we obtain by using (2.18) and (2.25),

$$\int_0^t \langle \bar{u}_t, \xi \rangle \, ds = (\bar{u}(\cdot, t), \xi) - (u_0, \xi),$$

which can be written as  $(\bar{u}(\cdot, t), \xi) - (\bar{u}(\cdot, 0), \xi) = (\bar{u}(\cdot, t), \xi) - (u_0, \xi)$ . Therefore,

$$\bar{u}(\cdot, 0) = u_0 \quad \text{a.e. in } \Omega.$$

Now, we see that  $\bar{u}$  satisfies

$$\left\{ \begin{array}{l} \bar{u} \in L^2(0, T; H_0^1(\Omega)), \quad \bar{u}_t \in L^2(0, T; H^{-1}(\Omega)), \\ \bar{u}(\cdot, 0) = u_0 \quad \text{a.e. in } \Omega, \\ \frac{d}{dt} (\bar{u}, \xi) + M(l(v_\infty)) \int_\Omega A(x, t, v_\infty) \nabla \bar{u} \cdot \nabla \xi \, dx = \int_\Omega g(x, t, v_\infty) \xi \, dx \\ \quad \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega), \end{array} \right.$$

and by uniqueness we have  $\bar{u} = u_\infty$ . In view of the above, we observe that every subsequence of  $(u_j)_{j \in \mathbb{N}}$  has a sub-subsequence that converges to the same limit  $u_\infty$ . So, the sequence  $(F(v_j))_{j \in \mathbb{N}}$  converges to  $F(v_\infty)$  in  $\overline{B}(0, R_4)$ , and then,  $F: \overline{B}(0, R_4) \rightarrow \overline{B}(0, R_4)$  is continuous. Thus, by the Schauder fixed point theorem, there exists a fixed point  $u$  of  $F$  in  $E$  which is a solution of (1.6) since  $E \subset C([0, T]; L^2(\Omega))$ .  $\square$



### 3. UNIQUENESS AND CONTINUITY RESULTS OF SOLUTIONS

In this section, we assume that  $A(x, t, u) = a(x, t)$  depends only on the variable  $(x, t)$ . We investigate two uniqueness theorems of solutions for (1.6) and give a continuity result depending on the initial data assuming that  $M$  is a Lipschitz continuous function,

$$(3.1) \quad \exists m_2 \geq 0: |M(s_1) - M(s_2)| \leq m_2 |s_1 - s_2| \quad \forall s_1, s_2 \in \mathbb{R}.$$

In the following first uniqueness theorem, we are concerned with the case  $p = 1$  in which the function  $g$  satisfies a generalized Lipschitz condition.

**Theorem 3.1.** *Assume that (1.1)–(1.3), (3.1) hold and that  $g$  satisfies (1.4) with  $p = 1$ . Then the solution of (1.6) is unique.*

**PROOF.** Let  $u_1$  and  $u_2$  be two solutions of (1.6) corresponding to the same initial data  $u_0 \in L^2(\Omega)$ . Then for all  $\xi \in H_0^1(\Omega)$  we have

$$\begin{aligned} \frac{d}{dt}(u_1, \xi) + M(l(u_1)) \int_{\Omega} a(x, t) \nabla u_1 \cdot \nabla \xi \, dx &= \int_{\Omega} g(x, t, u_1) \xi \, dx, \\ \frac{d}{dt}(u_2, \xi) + M(l(u_2)) \int_{\Omega} a(x, t) \nabla u_2 \cdot \nabla \xi \, dx &= \int_{\Omega} g(x, t, u_2) \xi \, dx \end{aligned}$$

in the distributional sense in  $\mathcal{D}'(0, T)$ . Choosing  $\xi = u_1 - u_2$  and subtracting the two equations from each other, we arrive at

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_1 - u_2\|_{L^2(\Omega)}^2 + M(l(u_1)) \int_{\Omega} a(x, t) \nabla u_1 \cdot \nabla (u_1 - u_2) \, dx \\ - M(l(u_2)) \int_{\Omega} a(x, t) \nabla u_2 \cdot \nabla (u_1 - u_2) \, dx \\ = \int_{\Omega} (g(x, t, u_1) - g(x, t, u_2))(u_1 - u_2) \, dx, \end{aligned}$$

which can be written as

$$(3.2) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_1 - u_2\|_{L^2(\Omega)}^2 + M(l(u_1)) \int_{\Omega} a(x, t) \nabla (u_1 - u_2) \cdot \nabla (u_1 - u_2) \, dx \\ = (M(l(u_2)) - M(l(u_1))) \int_{\Omega} a(x, t) \nabla u_2 \cdot \nabla (u_1 - u_2) \, dx \\ + \int_{\Omega} (g(x, t, u_1) - g(x, t, u_2))(u_1 - u_2) \, dx. \end{aligned}$$

Using (1.1)–(1.3), (1.4) for  $p = 1$ , (3.1) and Cauchy-Schwarz inequality, we obtain from (3.2),

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u_1 - u_2\|_{L^2(\Omega)}^2 + \lambda m_0 \int_{\Omega} |\nabla(u_1 - u_2)|^2 dx \\ & \leq m_2 \Lambda |l(u_2) - l(u_1)| \int_{\Omega} |\nabla u_2| |\nabla(u_1 - u_2)| dx + \int_{\Omega} |b_1(x, t)| |u_1 - u_2|^2 dx \\ & \leq m_2 \Lambda \|\phi\|_{L^2(\Omega)} \|u_1 - u_2\|_{L^2(\Omega)} \|u_2\|_{H_0^1(\Omega)} \|u_1 - u_2\|_{H_0^1(\Omega)} \\ & \quad + \|b_1\|_{L^\infty(Q)} \|u_1 - u_2\|_{L^2(\Omega)}^2, \end{aligned}$$

which, by using Young's inequality, leads to

$$\begin{aligned} (3.3) \quad & \frac{1}{2} \frac{d}{dt} \|u_1 - u_2\|_{L^2(\Omega)}^2 + \lambda m_0 \|u_1 - u_2\|_{H_0^1(\Omega)}^2 \\ & \leq \left\{ \frac{(m_2 \Lambda \|\phi\|_{L^2(\Omega)} \|u_2\|_{H_0^1(\Omega)})^2}{2\lambda m_0} + \|b_1\|_{L^\infty(Q)} \right\} \|u_1 - u_2\|_{L^2(\Omega)}^2 \\ & \quad + \frac{\lambda m_0}{2} \|u_1 - u_2\|_{H_0^1(\Omega)}^2. \end{aligned}$$

From (3.3), we obtain for a function  $\theta \in L^1(0, T)$ ,

$$(3.4) \quad \frac{d}{dt} \|u_1 - u_2\|_{L^2(\Omega)}^2 \leq \theta(t) \|u_1 - u_2\|_{L^2(\Omega)}^2.$$

Multiplying (3.4) by  $\exp(-\int_0^t \theta(s) ds)$ , we get

$$\frac{d}{dt} \left( \exp\left(-\int_0^t \theta(s) ds\right) \|u_1 - u_2\|_{L^2(\Omega)}^2 \right) \leq 0.$$

Finally, integrating from 0 to  $t$  and taking into account that  $u_1(\cdot, 0) = u_2(\cdot, 0)$  a.e. in  $\Omega$ , we find  $u_1 = u_2$  a.e. in  $Q$ .  $\square$

With the assumption that the function  $g$  is decreasing with respect to  $u$ , we state and prove our second uniqueness theorem.

**Theorem 3.2.** *Assume that (1.1)–(1.3), (3.1) and either (1.4) (with  $p \in (0, 1)$ ) or (1.5) hold. In addition, for a.e.  $(x, t) \in Q$ ,*

$$(3.5) \quad \text{the function } s \mapsto g(x, t, s) \text{ is decreasing on } \mathbb{R}.$$

*Then the solution of (1.6) is unique.*

Proof. Let  $u_1$  and  $u_2$  be two solutions of (1.6) corresponding to the same initial data  $u_0 \in L^2(\Omega)$ . Setting  $w = u_1 - u_2$ , we can see that the function  $w$  satisfies

$$(3.6) \quad \begin{cases} \frac{d}{dt}(w, \xi) + M(l(u_1)) \int_{\Omega} a(x, t) \nabla w \cdot \nabla \xi \, dx + \int_{\Omega} G(x, t) \xi \, dx \\ \quad = (M(l(u_2)) - M(l(u_1))) \int_{\Omega} a(x, t) \nabla u_2 \cdot \nabla \xi \, dx + \int_{\Omega} w \xi \, dx, \\ \quad \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega), \\ w(\cdot, 0) = 0 \quad \text{a.e. in } \Omega, \end{cases}$$

where  $G(x, t) = g(x, t, u_2) - g(x, t, u_1) + w$ . Choosing  $\xi = w$  as test function in the first equation of (3.6) and integrating from 0 to  $t$ , we obtain

$$(3.7) \quad \begin{aligned} \frac{1}{2} \|w\|_{L^2(\Omega)}^2 + \int_0^t M(l(u_1)) \int_{\Omega} a(x, s) \nabla w \cdot \nabla w \, dx \, ds + \int_0^t \int_{\Omega} G(x, s) w \, dx \, ds \\ = \int_0^t (M(l(u_2)) - M(l(u_1))) \int_{\Omega} a(x, s) \nabla u_2 \cdot \nabla w \, dx \, ds + \int_0^t \|w\|_{L^2(\Omega)}^2 \, ds. \end{aligned}$$

Using (1.1)–(1.3), (3.1), (3.5), the Cauchy-Schwarz inequality and Young's inequality, we get

$$(3.8) \quad \int_0^t M(l(u_1)) \int_{\Omega} a(x, s) \nabla w \cdot \nabla w \, dx \, ds \geq m_0 \lambda \int_0^t \|w\|_{H_0^1(\Omega)}^2 \, ds,$$

$$(3.9) \quad \begin{aligned} \left| \int_0^t (M(l(u_2)) - M(l(u_1))) \int_{\Omega} a(x, s) \nabla u_2 \cdot \nabla w \, dx \, ds \right| \\ \leq m_2 \Lambda \int_0^t \int_{\Omega} |\phi| |w| \, dx \int_{\Omega} |\nabla u_2| |\nabla w| \, dx \, ds \\ \leq m_2 \Lambda \|\phi\|_{L^2(\Omega)} \int_0^t \|w\|_{L^2(\Omega)} \|u_2\|_{H_0^1(\Omega)} \|w\|_{H_0^1(\Omega)} \, ds \\ \leq \frac{m_0 \lambda}{2} \int_0^t \|w\|_{H_0^1(\Omega)}^2 \, ds \\ \quad + \frac{(m_2 \Lambda \|\phi\|_{L^2(\Omega)})^2}{2m_0 \lambda} \int_0^t \|w\|_{L^2(\Omega)}^2 \|u_2\|_{H_0^1(\Omega)}^2 \, ds \end{aligned}$$

and

$$(3.10) \quad \int_0^t \int_{\Omega} G(x, s) w \, dx \, ds = \int_0^t \int_{\Omega} (g(x, s, u_2) - g(x, s, u_1) + w) w \, dx \, ds \geq 0.$$

Combining (3.8)–(3.10), we obtain from (3.7),

$$\begin{aligned} \frac{1}{2} \|w\|_{L^2(\Omega)}^2 + \frac{m_0 \lambda}{2} \int_0^t \|w\|_{H_0^1(\Omega)}^2 \, ds \\ \leq \frac{(m_2 \Lambda \|\phi\|_{L^2(\Omega)})^2}{2m_0 \lambda} \int_0^t \|w\|_{L^2(\Omega)}^2 \|u_2\|_{H_0^1(\Omega)}^2 \, ds + \int_0^t \|w\|_{L^2(\Omega)}^2 \, ds, \end{aligned}$$

which leads to

$$\|w\|_{L^2(\Omega)}^2 \leq \int_0^t \left\{ \frac{(m_2 \Lambda \|\phi\|_{L^2(\Omega)})^2}{m_0 \lambda} \|u_2\|_{H_0^1(\Omega)}^2 + 2 \right\} \|w\|_{L^2(\Omega)}^2 ds.$$

From Lemma 3.1 of [14] we have  $\|w\|_{L^2(\Omega)} = 0$  in  $[0, T]$ , which implies that  $w = 0$  a.e. in  $Q$ , and then,  $u_1 = u_2$  a.e. in  $Q$ .  $\square$

Our continuity result is now presented.

**Theorem 3.3.** *Suppose either Theorem 3.1 or 3.2 holds. Then for all  $t \in [0, T]$ , the mapping*

$$\mathcal{T}: L^2(\Omega) \rightarrow L^2(\Omega), \quad u_0 \mapsto u(\cdot, t),$$

where  $u$  is the unique solution of (1.6) corresponding to the initial data  $u_0$ , satisfies the following continuity result:

$$u_0^j \rightharpoonup u_0 \text{ weakly in } L^2(\Omega) \Rightarrow \mathcal{T}(u_0^j) \rightarrow \mathcal{T}(u_0) \text{ strongly in } L^2(\Omega).$$

Particularly,  $\mathcal{T}$  is weakly continuous.

*Proof.* Let  $(u_0^j)_{j \in \mathbb{N}} \subset L^2(\Omega)$  be a sequence of initial data which converges to  $u_0$  weakly in  $L^2(\Omega)$ , and let  $u^j, u$  be, respectively, the solutions of (1.6) corresponding to  $u_0^j, u_0, j \in \mathbb{N}$ . For all  $j \in \mathbb{N}$  and all  $p \in (0, 1)$  we have

$$(3.11) \quad \|u^j\|_{L^2(Q)} \leq T^{1/2} \left\{ C_\Omega \left( \frac{2}{\lambda m_0} \right)^{1/2} (\|b_p\|_{L^{2/(1-p)}(Q)} \|u^j\|_{L^2(Q)}^p + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) \right. \\ \left. + \|u_0^j\|_{L^2(\Omega)} \right\},$$

$$(3.12) \quad \|u^j\|_{L^2(\Omega)} \leq \left\{ \frac{2C_\Omega^2}{\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)}^2 \|u^j\|_{L^2(Q)}^{2p} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) \right. \\ \left. + \|u_0^j\|_{L^2(\Omega)}^2 \right\}^{1/2}$$

and

$$(3.13) \quad \|u^j\|_{L^2(0, T; H_0^1(\Omega))} \leq \frac{1}{(\lambda m_0)^{1/2}} \left\{ \frac{2C_\Omega^2}{\lambda m_0} (\|b_p\|_{L^{2/(1-p)}(Q)}^2 \|u^j\|_{L^2(Q)}^{2p} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) \right. \\ \left. + \|u_0^j\|_{L^2(\Omega)}^2 \right\}^{1/2},$$

and if  $p = 1$ ,

$$(3.14) \quad \|u^j\|_{L^2(Q)} \leq T^{1/2} \left\{ C_\Omega \left( \frac{2}{\lambda m_0} \right)^{1/2} (\|b_1\|_{L^\infty(Q)} \|u^j\|_{L^2(Q)} + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}) \right. \\ \left. + \|u_0^j\|_{L^2(\Omega)} \right\},$$

$$(3.15) \quad \|u^j\|_{L^2(\Omega)} \leq \left\{ \frac{2C_\Omega^2}{\lambda m_0} (\|b_1\|_{L^\infty(Q)}^2 \|u^j\|_{L^2(Q)}^2 + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \|u_0^j\|_{L^2(\Omega)}^2 \right\}^{1/2}$$

and

$$(3.16) \quad \|u^j\|_{L^2(0,T;H_0^1(\Omega))} \leq \frac{1}{(\lambda m_0)^{1/2}} \left\{ \frac{2C_\Omega^2}{\lambda m_0} (\|b_1\|_{L^\infty(Q)}^2 \|u^j\|_{L^2(Q)}^2 + \|g(\cdot, \cdot, 0)\|_{L^2(Q)}^2) + \|u_0^j\|_{L^2(\Omega)}^2 \right\}^{1/2}.$$

On the other hand, for all  $\xi \in \mathcal{D}(0, T; H_0^1(\Omega))$ ,

$$(3.17) \quad \int_Q u^j \xi_t \, dx \, dt = \int_0^T M(l(u^j)) \int_\Omega a(x, t) \nabla u^j \cdot \nabla \xi \, dx \, dt - \int_Q g(x, t, u^j) \xi \, dx \, dt.$$

Observe that the sequence  $(u_0^j)_{j \in \mathbb{N}}$  is bounded in  $L^2(\Omega)$  since it is a weakly convergent sequence in  $L^2(\Omega)$ . Then, by taking into account that  $C_\Omega(2T/(\lambda m_0))^{1/2} \|b_1\|_{L^\infty(Q)} < 1$  (in the case  $p = 1$ ), we deduce from (3.11) and (3.14) that  $(u^j)_{j \in \mathbb{N}}$  is bounded in  $L^2(Q)$ . Hence, using (3.12), (3.13) and (3.15)–(3.17), we can find a constant  $c_3 = c_3(p)$  independent of  $j$  such that for all  $j \in \mathbb{N}$ ,

$$(3.18) \quad \|u^j\|_{L^\infty(0,T;L^2(\Omega))} \leq c_3,$$

$$(3.19) \quad \|u^j\|_{L^2(0,T;H_0^1(\Omega))} \leq c_3,$$

$$(3.20) \quad \|u_t^j\|_{L^2(0,T;H^{-1}(\Omega))} \leq c_3.$$

Similarly, if  $g$  satisfies (1.5), we arrive at

$$\forall j \in \mathbb{N}: \|u^j\|_{L^2(\Omega)}^2 + \lambda m_0 \int_0^t \|u^j\|_{H_0^1(\Omega)}^2 \, ds \leq \frac{(C_\Omega \|h\|_{L^2(Q)})^2}{\lambda m_0} + \|u_0^j\|_{L^2(\Omega)}^2,$$

which leads us to saying that inequalities (3.18)–(3.20) are satisfied for another constant independent of  $j$ . Therefore, there exists a subsequence  $j_k$  and  $\bar{u} \in E$  such that

$$(3.21) \quad u^{j_k} \rightarrow \bar{u} \quad \text{strongly in } L^2(Q),$$

$$(3.22) \quad u^{j_k} \rightarrow \bar{u} \quad \text{a.e. in } Q,$$

$$(3.23) \quad M(l(u^{j_k})) \rightarrow M(l(\bar{u})) \quad \text{strongly in } L^2(0, T),$$

$$(3.24) \quad u^{j_k}(\cdot, t) \rightarrow \bar{u}(\cdot, t) \quad \text{strongly in } L^2(\Omega), \text{ a.e. } t \in (0, T),$$

$$(3.25) \quad u^{j_k} \rightharpoonup \bar{u} \quad \text{weakly in } L^2(0, T; H_0^1(\Omega)),$$

$$(3.26) \quad u_t^{j_k} \rightharpoonup \bar{u}_t \quad \text{weakly in } L^2(0, T; H^{-1}(\Omega)),$$

$$(3.27) \quad u^{j_k} \rightharpoonup^* \bar{u} \quad \text{weakly star in } L^\infty(0, T; L^2(\Omega)),$$

$$(3.28) \quad u_0^{j_k} \rightarrow u_0 \quad \text{weakly in } L^2(\Omega).$$

We have

$$(3.29) \quad \int_0^T \varphi(t) M(l(u^{j_k})) \int_{\Omega} a(x, t) \nabla u^{j_k} \cdot \nabla \xi \, dx \, dt - \int_Q u^{j_k} \xi \varphi_t \, dx \, dt \\ = \int_Q g(x, t, u^{j_k}) \xi \varphi \, dx \, dt \quad \forall k \in \mathbb{N}, \quad \forall \varphi \in \mathcal{D}(0, T), \quad \forall \xi \in H_0^1(\Omega).$$

We notice that if one of assumptions (1.4) and (1.5) holds, we obtain

$$(3.30) \quad \lim_{k \rightarrow \infty} \int_Q g(x, t, u^{j_k}) \xi \varphi \, dx \, dt = \int_Q g(x, t, \bar{u}) \xi \varphi \, dx \, dt$$

by using, respectively, (3.21) and (3.22), then, letting  $k \rightarrow \infty$  in (3.29) we get by taking into account (1.2), (3.21), (3.23), (3.25) and (3.30),

$$\int_0^T \varphi(t) M(l(\bar{u})) \int_{\Omega} a(x, t) \nabla \bar{u} \cdot \nabla \xi \, dx \, dt - \int_Q \bar{u} \xi \varphi_t \, dx \, dt = \int_Q g(x, t, \bar{u}) \xi \varphi \, dx \, dt.$$

So, the limit  $\bar{u}$  is a solution to

$$\begin{cases} \bar{u} \in L^2(0, T; H_0^1(\Omega)) \cap C([0, T]; L^2(\Omega)), & \bar{u}_t \in L^2(0, T; H^{-1}(\Omega)), \\ \frac{d}{dt}(\bar{u}, \xi) + M(l(\bar{u})) \int_{\Omega} a(x, t) \nabla \bar{u} \cdot \nabla \xi \, dx = \int_{\Omega} g(x, t, \bar{u}) \xi \, dx \\ \text{in } \mathcal{D}'(0, T) \quad \forall \xi \in H_0^1(\Omega). \end{cases}$$

Moreover,

$$\forall \xi \in H_0^1(\Omega), \quad \text{a.e. } t \in (0, T): \quad \int_0^t \langle u_t^{j_k}, \xi \rangle \, ds = (u^{j_k}(\cdot, t), \xi) - (u_0^{j_k}, \xi).$$

Passing to the limit as  $k \rightarrow \infty$  and using (3.24), (3.26) and (3.28),

$$\int_0^t \langle \bar{u}_t, \xi \rangle \, ds = (\bar{u}(\cdot, t), \xi) - (u_0, \xi),$$

which leads to

$$(\bar{u}(\cdot, t), \xi) - (\bar{u}(\cdot, 0), \xi) = (\bar{u}(\cdot, t), \xi) - (u_0, \xi).$$

Hence,  $\bar{u}(\cdot, 0) = u_0$  a.e. in  $\Omega$ . Therefore, due to the uniqueness of the limit, we have  $\bar{u} = u$  a.e. in  $Q$ . Thus, by (3.27),

$$u^{j_k} \rightharpoonup^* u \quad \text{weakly star in } L^\infty(0, T; L^2(\Omega)).$$

Particularly, if  $\xi \in H_0^1(\Omega)$ , it follows that

$$(3.31) \quad (u^{j_k}(\cdot, t), \xi) \rightharpoonup^* (u(\cdot, t), \xi) \quad \text{weakly star in } L^\infty(0, T).$$

Now, let  $t_1$  and  $t_2$  be elements of  $[0, T]$  such that  $t_2 > t_1$ . We have

$$\begin{aligned}
 (3.32) \quad (u^{j_k}(\cdot, t_2), \xi) - (u^{j_k}(\cdot, t_1), \xi) &= \int_{t_1}^{t_2} \langle u_t^{j_k}, \xi \rangle dt \leq \int_{t_1}^{t_2} \|u_t^{j_k}\|_{H^{-1}(\Omega)} \|\xi\|_{H_0^1(\Omega)} dt \\
 &\leq (t_2 - t_1)^{1/2} \|\xi\|_{H_0^1(\Omega)} \|u_t^{j_k}\|_{L^2(0, T; H^{-1}(\Omega))} \\
 &\leq c \|\xi\|_{H_0^1(\Omega)} (t_2 - t_1)^{1/2}.
 \end{aligned}$$

Due to (3.32), the sequence of functions  $(u^{j_k}(\cdot, t), \xi)$  is equicontinuous, and then, it is relatively compact in  $C([0, T])$ . By (3.31) and the uniqueness of the limit, we have up to a subsequence

$$\forall \xi \in H_0^1(\Omega): (u^{j_k}(\cdot, t), \xi) \rightarrow (u(\cdot, t), \xi) \quad \text{in } C([0, T]).$$

Thanks to the density of  $H_0^1(\Omega)$  in  $L^2(\Omega)$ ,

$$\forall \xi \in L^2(\Omega): (u^{j_k}(\cdot, t), \xi) \rightarrow (u(\cdot, t), \xi) \quad \text{strongly in } \mathbb{R} \quad \forall t \in [0, T],$$

which means that

$$u^{j_k}(\cdot, t) \rightarrow u(\cdot, t) \quad \text{strongly in } L^2(\Omega) \quad \forall t \in [0, T].$$

Since every subsequence of  $(u^j(\cdot, t))_{j \in \mathbb{N}}$  has a sub-subsequence that converges to the same limit  $u(\cdot, t)$ , we deduce that

$$\mathcal{T}(u_0^j) = u^j(\cdot, t) \rightarrow \mathcal{T}(u_0) = u(\cdot, t) \quad \text{strongly in } L^2(\Omega) \quad \forall t \in [0, T],$$

which completes the proof of the theorem. □

### References

- [1] *C. O. Alves, T. Boudjeriou*: Existence of solution for a class of nonlocal problem via dynamical methods. *Rend. Circ. Mat., Palermo (2)* 71 (2022), 611–632. [zbl](#) [MR](#) [doi](#)
- [2] *C. O. Alves, D.-P. Covei*: Existence of solution for a class of nonlocal elliptic problem via sub-supersolution method. *Nonlinear Anal., Real World Appl.* 23 (2015), 1–8. [zbl](#) [MR](#) [doi](#)
- [3] *C. T. Anh, L. T. Tinh, V. M. Toi*: Global attractors for nonlocal parabolic equations with a new class of nonlinearities. *J. Korean Math. Soc.* 55 (2018), 531–551. [zbl](#) [MR](#) [doi](#)
- [4] *D. Arcoya, T. Leonori, A. Primo*: Existence of solutions for semilinear nonlocal elliptic problems via a Bolzano theorem. *Acta Appl. Math.* 127 (2013), 87–104. [zbl](#) [MR](#) [doi](#)
- [5] *M. Bousselsal, E. Zaouche*: Existence of solution for nonlocal heterogeneous elliptic problems. *Mediterr. J. Math.* 17 (2020), Article ID 129, 10 pages. [zbl](#) [MR](#) [doi](#)
- [6] *M. Chipot*: *Elements of Nonlinear Analysis*. Birkhäuser Advanced Texts. Birkhäuser, Basel, 2000. [zbl](#) [MR](#) [doi](#)

- [7] *M. Chipot, B. Lovat*: Some remarks on non local elliptic and parabolic problems. *Nonlinear Anal., Theory Methods Appl.* *30* (1997), 4619–4627. [zbl](#) [MR](#) [doi](#)
- [8] *M. Chipot, B. Lovat*: On the asymptotic behaviour of some nonlocal problems. *Positivity* *3* (1999), 65–81. [zbl](#) [MR](#) [doi](#)
- [9] *M. Chipot, B. Lovat*: Existence and uniqueness results for a class of nonlocal elliptic and parabolic problems. *Dyn. Contin. Discrete Impuls. Syst., Ser. A, Math. Anal.* *8* (2001), 35–51. [zbl](#) [MR](#)
- [10] *M. Chipot, L. Molinet*: Asymptotic behavior of some nonlocal diffusion problems. *Appl. Anal.* *80* (2001), 279–315. [zbl](#) [MR](#) [doi](#)
- [11] *F. J. S. A. Corrêa*: On positive solutions of nonlocal and nonvariational elliptic problems. *Nonlinear Anal., Theory Methods Appl., Ser. A* *59* (2004), 1147–1155. [zbl](#) [MR](#) [doi](#)
- [12] *R. Dautray, J.-L. Lions*: *Mathematical Analysis and Numerical Methods for Science and Technology*. Vol. 5. *Evolution Problems I*. Springer, Berlin, 1992. [zbl](#) [MR](#)
- [13] *S. B. de Menezes*: Remarks on weak solutions for a nonlocal parabolic problem. *Int. J. Math. Math. Sci.* *2006* (2006), Article ID 82654, 10 pages. [zbl](#) [MR](#) [doi](#)
- [14] *J. K. Hale, S. M. Verduyn Lunel*: *Introduction to Functional Differential Equations*. *Applied Mathematical Sciences* 99. Springer, New York, 1993. [zbl](#) [MR](#) [doi](#)
- [15] *R. Jiang, C. Zhai*: Properties of unique positive solutions for a class of nonlocal semilinear elliptic equations. *Topol. Methods Nonlinear Anal.* *50* (2017), 669–682. [zbl](#) [MR](#) [doi](#)
- [16] *F. Jin, B. Yan*: Existence and global behavior of the solution to a parabolic equation with nonlocal diffusion. *AIMS Math.* *6* (2021), 5292–5315. [zbl](#) [MR](#) [doi](#)
- [17] *B. Lovat*: *Études de quelques problèmes paraboliques non locaux*: Thèse. Université de Metz, Metz, 1995 (In French.); Available at <https://hal.univ-lorraine.fr/tel-01777092>.
- [18] *B. Yan, D. Wang*: The multiplicity of positive solutions for a class of nonlocal elliptic problem. *J. Math. Anal. Appl.* *442* (2016), 72–102. [zbl](#) [MR](#) [doi](#)
- [19] *E. Zaouche*: Nontrivial weak solutions for nonlocal nonhomogeneous elliptic problems. *Appl. Anal.* *101* (2022), 1261–1270. [zbl](#) [MR](#) [doi](#)
- [20] *E. Zaouche*: Existence theorems of nontrivial and positive solutions for nonlocal inhomogeneous elliptic problems. *Ric. Mat.* *72* (2023), 949–960. [zbl](#) [MR](#) [doi](#)

*Authors' addresses*: *Tayeb Benhamoud, Elmehdi Zaouche* (corresponding author), Department of Mathematics, University of El Oued, El Oued 39000, Algeria, e-mail: [tayeb06@yahoo.fr](mailto:tayeb06@yahoo.fr), [elmehdi-zaouche@univ-eloued.dz](mailto:elmehdi-zaouche@univ-eloued.dz), [elmehdizaouche45@gmail.com](mailto:elmehdizaouche45@gmail.com); *Mahmoud Bousselsal*, Department of Mathematics, Laboratoire des équations aux dérivées partielles et H.M., École Normale Supérieure, Vieux-Kouba, Algiers, Algeria, e-mail: [bousselsal155@gmail.com](mailto:bousselsal155@gmail.com).